



CONCUSSION INFORMATION SHEET

A concussion is a brain injury and all brain injuries are serious. They are caused by a bump, blow, or jolt to the head, or by a blow to another part of the body with the force transmitted to the head. They can range from mild to severe and can disrupt the way the brain normally works. Even though most concussions are mild, **all concussions are potentially serious and may result in complications including prolonged brain damage and death if not recognized and managed properly.** and other

- Nausea or vomiting
- Neck pain
- Balance problems or dizziness
- Blurred, double or fuzzy vision
- Sensitivity to light or noise
- Feeling sluggish or slowed down
- Feeling foggy or groggy

- Don't feel right
- Fatigue or low energy
- Sadness
- Nervousness or anxiety
- Irritability
- More emotional
- Confusion
-



What can happen if my child keeps on playing with a concussion or returns too soon

Athletes with the signs and symptoms of concussion should be removed from play immediately. Continuing to play with the signs and symptoms of a concussion leaves the young athlete especially vulnerable to greater injury. There is an increased risk of significant damage from a concussion for a period of time after the concussion occurs, particularly if the athlete suffers another concussion before completely recovering from the first one. This can lead to prolonged recovery or even to severe brain swelling (second impact syndrome) with devastating and even fatal consequences. It is well known that